

Journal Pre-proof

AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis

Joseph D. Feuerstein, MD, Kim L. Isaacs, MD, Yechezkel Schneider, MD, Shazia Mehmood Siddique, MD, MS, Yngve Falck-Ytter, MD, Siddharth Singh, MD, MS, on behalf of the American Gastroenterological Association Institute Clinical Guidelines Committee

PII: S0016-5085(20)30018-4
DOI: <https://doi.org/10.1053/j.gastro.2020.01.006>
Reference: YGAST 63119

To appear in: *Gastroenterology*

Please cite this article as: Feuerstein JD, Isaacs KL, Schneider Y, Siddique SM, Falck-Ytter Y, Singh S, on behalf of the American Gastroenterological Association Institute Clinical Guidelines Committee, AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis, *Gastroenterology* (2020), doi: <https://doi.org/10.1053/j.gastro.2020.01.006>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 by the AGA Institute



AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis

Joseph D. Feuerstein MD¹, Kim L. Isaacs MD², Yecheskel Schneider MD³, Shazia Mehmood Siddique MD, MS³, Yngve Falck-Ytter MD⁴, and Siddharth Singh MD, MS⁵ on behalf of the American Gastroenterological Association Institute Clinical Guidelines Committee

¹ Division of Gastroenterology and Center for Inflammatory Bowel Diseases, Beth Israel Deaconess Medical Center, Boston, MA

³ Division of Gastroenterology, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA

⁴ Division of Gastroenterology, Case Western Reserve University, and VA Northeast Ohio Healthcare System, Cleveland, Ohio

⁵ Division of Gastroenterology, University of California, San Diego, La Jolla, California

Acknowledgements: The AGA Clinical Guidelines Committee included Karen Chachu MD, PhD, Lukejohn Day MD, AGAF, Benjamin Lebwohl MD, AGAF, Thiruvengadam Muniraj MD, PhD, MRCP, Amit Patel MD, Anne F. Peery, Raj Shah, Shahnaz Sultan, Harminder Singh, Siddharth Singh, Stuart Spechler, Grace Su, Aaron P. Thrift, Jennifer M. Weiss, and Adam V. Weizman

Address for Correspondence:

Chair, Clinical Guidelines Committee,
American Gastroenterological Association
National Office, 4930 Del Ray Avenue,
Bethesda, Maryland 20814
E-mail: clinicalpractice@gastro.org
Telephone: (301) 941-2618

This document presents the official recommendations of the American Gastroenterological Association (AGA) on the management of moderate to severe ulcerative colitis. The guideline was developed by the AGA's Clinical Practice Guideline Committee and approved by the AGA Governing Board. Development of this guideline and its accompanying technical review was fully funded by the AGA Institute with no additional outside funding.

Conflict of interest disclosure: All members were required to complete the disclosure statement. These statements are maintained at the American Gastroenterological Association (AGA) headquarters in Bethesda, Maryland, and pertinent disclosures are published with this report.

Conflicts of interest:

Joseph D. Feuerstein has no relevant conflict to report.

Kim L. Isaacs received research support for multi-center clinical trials and safety registries from Abbvie, Takeda, UCB, Janssen, and Hoffman-Laroche.

Yechezkel Schneider has no relevant conflict to report.

Shazia Mehmood Siddique has no relevant conflict to report.

Yngve Falck-Ytter MD has no relevant conflict to report.

Siddharth Singh MD has no relevant conflict to report.

Anticipated update: 3 years from publication (2023)

Guideline Panel: Joseph Feuerstein (Chair), Siddharth Singh (Co-chair, GRADE Methodologist), Kim Isaacs (Member), Yechezkel Schneider (Member), Yngve Falck-Ytter (Member, Chair, Clinical Guidelines Committee), Shazia Mehmood Siddique (Trainee GRADE Methodologist)

Technical Review Panel: Jessica Allegretti (Content expert), Jonathan Terdiman (Content expert), Siddharth Singh (GRADE Methodologist), Shazia Mehmood Siddique (Trainee GRADE Methodologist)

INTRODUCTION

This document presents the official recommendations of the American Gastroenterological Association (AGA) on the management of moderate to severe ulcerative colitis (UC). The guideline was developed by the AGA Institute's Clinical Guidelines Committee and approved by the AGA Governing Board. It is accompanied by a technical review that provides a detailed synthesis of the evidence from which these recommendations were formulated.¹ Development of this guideline and the accompanying technical review was fully funded by the AGA Institute without additional outside funding. Members of the guideline panel and technical review panel were selected by the AGA Governing Board in consultation with the Clinical Guidelines Committee with careful consideration of all Institute of Medicine recommendations for clinical guideline development. Joseph Feuerstein was the guideline panel chair and Siddharth Singh was the methodologist and co-chair of the guideline panel. A patient representative was also included in the development and review process and had no recommended changes. The guideline and accompanying technical review underwent independent peer review, and a 30-day open public comment period; all comments were collated by the AGA staff, and were reviewed and carefully considered by the guideline panel and technical review teams, respectively. Changes were incorporated in revised documents, and where changes were not accepted, a thoughtful response document was created. Following the public comment period, two pivotal clinical trials (VARSITY, UNIFI) were published and a critical safety update on tofacitinib was issued by the Food and Drug Administration (FDA). At the recommendation of the Clinical Guidelines Committee, the technical review and clinical guidelines were updated to

incorporate this new evidence as presented here. In accordance with the Clinical Guidelines Committee policies, all clinical guidelines are reviewed annually at the AGA Clinical Guideline Committee meeting for new information. The next update for these guidelines is anticipated in three years from publication.

UC is a chronic inflammatory bowel disease with peak onset in early adulthood.² Untreated, the natural history of the disease is one of relapsing and remitting mucosal inflammation. Based on population-based cohort studies, the majority of patients with UC have a mild to moderate course, generally most active at diagnosis and then in varying periods of remission or mild activity. Approximately 15% patients may experience an aggressive course, and 20% of these patients may require hospitalization for severe disease activity.^{3, 4} The 5- and 10-year cumulative risk of colectomy is 10-15%, primarily limited to patients with moderate to severe disease activity; a subset of hospitalized patients with acute severe ulcerative colitis (ASUC) have short-term colectomy rates of 25-30%.⁴ Predictors of an aggressive disease course and colectomy are: young age at diagnosis (age < 40 years old), extensive disease, severe endoscopic activity (presence of large and/or deep ulcers), presence of extra-intestinal manifestations, early need for corticosteroids and elevated inflammatory markers.⁵ For this guideline and the accompanying technical review, moderate to severe UC is defined based on the Truelove and Witts criteria and Mayo Clinic score (Table 1).⁵⁻⁷ After excluding concomitant infections (such as *Clostridium difficile*), patients with moderate to severe disease are those who are dependent on or refractory to corticosteroids, have severe endoscopic disease activity (presence of ulcers), or are at high risk of colectomy. When reported, Mayo Clinic scores of 6-12 with an endoscopic subscore of 2 or 3 were considered moderate to severe disease. ASUC in this guideline is defined as hospitalized patients with the following Truelove and Witts criteria: ≥ 6 bloody bowel movements/day with at least one marker of systemic toxicity including heart rate >90 beats/minute, temperature >37.8⁰ C, hemoglobin <10.5 g/dl and/or erythrocyte sedimentation rate >30 mm/h.⁷

There are a number of different drug classes for long-term management of moderate to severe UC, including tumor necrosis factor (TNF)- α antagonists, anti-integrin agent (vedolizumab), janus kinase inhibitor (tofacitinib) and immunomodulators (thiopurines, methotrexate).⁸ In general, most drugs that are initiated for induction of remission are continued as maintenance therapy, if they are effective. This clinical practice is considered standard of care in this guideline and it is assumed that if a drug (excluding corticosteroids and cyclosporine) is started for and is effective for induction of remission or response, it will be continued for maintenance of remission.

This guideline addresses the medical management of adult outpatients with moderate to severe UC as well as the medical management of adult hospitalized patients with ASUC. The guideline focuses on immunomodulators, biologics, and small molecules for induction and maintenance of remission (for moderate to severe UC) and decreasing the risk of colectomy (for ASUC). As noted above, unless otherwise specified, we do not present separate recommendations for induction and maintenance of remission. The drugs are listed in order of FDA approval unless specifically mentioned. The first seven questions are focused on the medical management of adult outpatients with moderate to severe UC; the subsequent four questions are focused on adult patients hospitalized with ASUC, focusing on initial management, and rescue therapy in cases of corticosteroid-refractory disease. We acknowledge challenges in defining moderate disease activity and severity, with variable definition in clinical practice, and an understanding of this entity may be enhanced reading the AGA guideline on the management of mild to moderate UC.⁹ This guideline does not address surgical management of moderate to severe UC or ASUC. Therapeutic drug monitoring to guide the use of biologic therapy has been addressed in separate AGA guidelines.¹⁰ The guideline is intended for the use of gastroenterology providers, primary care providers, surgeons, patients and policymakers.

For this guideline, critical outcomes for decision-making for adults with moderate to severe UC were induction and maintenance of remission, and for hospitalized adults with ASUC was short-term colectomy risk (within 3 months of hospitalization), and are reported in the evidence profiles. Important outcomes of interest were induction and maintenance of endoscopic remission, maintenance of corticosteroid-free remission, serious adverse events (including serious infections and malignancy), and treatment tolerability (drug discontinuation due to adverse events). These were considered in evidence synthesis especially if inadequate or conflicting data was observed for critical outcomes. Safety considerations with these medications have been synthesized in the accompanying technical review. In the recommendations presented in this guideline, estimates of effects of different medications are presented as the risk for failure to induce or maintain remission, i.e., a relative risk (RR) or odds ratio (OR) <1 suggests that the drug under consideration is more effective than the comparison drug or placebo for induction or maintenance of remission.

This guideline was developed using a process described elsewhere.¹¹ Briefly, the AGA process for developing clinical practice guidelines incorporate GRADE methodology¹² and best practices as outlined by the Institute of Medicine.¹³ GRADE methodology was used to prepare the background information for the guideline and the accompanying technical review. Optimal understanding of the guideline will be enhanced by reading the applicable portions of the technical review. The guideline panel and the authors of the technical review met face to face on December 14, 2018 to discuss the findings from the technical review. The guideline authors subsequently formulated the guideline recommendations using the GRADE evidence-to-decision framework guidance. New evidence was presented to the guideline panel by the technical review team on October 16, 2019, and was reviewed and approved in a virtual face-to-face meeting on November 1, 2019. Although the quality of evidence (Table 2) was a key factor in determining the strength of the recommendations (Table 3), the panel also considered the balance between benefit and harm of interventions, patients' values and preferences, and

overall resource utilization. The recommendations, quality of evidence, and strength of recommendations are summarized in Table 4.

Journal Pre-proof

RECOMMENDATIONS

1. In adult outpatients with moderate-severe ulcerative colitis, the AGA recommends using infliximab, adalimumab, golimumab, vedolizumab, tofacitinib or ustekinumab over no treatment. (Strong recommendation, moderate quality evidence)

The panel recommends treating adult outpatients with moderate-severe UC with infliximab, adalimumab, golimumab, vedolizumab, tofacitinib or ustekinumab over no treatment for the induction and maintenance of remission. There were 16 randomized controlled trials (RCTs) comparing the TNF α antagonists, vedolizumab, tofacitinib and ustekinumab to placebo (see technical review). Induction of remission was assessed at 6-8 weeks and maintenance of remission was evaluated at 30-54 weeks. All active interventions were superior to placebo for induction of remission, regardless of prior biologic exposure (infliximab: RR, 2.85; 95% CI, 2.11-3.86; adalimumab: RR, 1.62; 95% CI, 1.15-2.29; golimumab RR, 2.49; 95% CI, 1.58-3.93; vedolizumab RR, 2.22; 95% CI, 1.36-3.64; tofacitinib RR, 3.22; 95% CI, 2.03-5.08; ustekinumab RR, 2.91; 95% CI, 1.72-4.94). Likewise, all active interventions were superior to placebo for maintenance of remission (infliximab: RR, 2.25; 95% CI, 1.67-3.05; adalimumab: RR, 2.28; 95% CI, 1.52-3.42; golimumab RR, 1.88; 95% CI, 1.32-2.68; vedolizumab RR, 2.31; 95% CI, 1.63-3.28; tofacitinib 5mg twice daily RR, 3.09; 95% CI, 1.99-4.79; ustekinumab RR, 1.83; 95% CI, 1.33-2.49). All medications were well-tolerated with low rates of serious adverse events in both trials of induction and maintenance therapy, not significantly different from placebo. Importantly, the recommended induction dose of tofacitinib is 10mg twice daily for 8 weeks; in select cases with modest response to initial 8 week therapy, high dose tofacitinib may considered for total 16 weeks. For long-term maintenance, tofacitinib 5mg twice daily is recommended for most patients; a higher dose may be considered in patients who lose response at 5mg twice daily dose after careful of risks and benefits of the medication. At higher doses, an unexpected increase in risk of pulmonary embolism and all-cause mortality has been observed.

The overall quality of evidence for this recommendation was moderate for both induction and maintenance of remission. While majority of the studies were registration trials sponsored by industry, there was no important bias, inconsistency or indirectness. The evidence was rated down for imprecision due to the lower number of events (< 200) for all comparisons, failing to achieve the optimal information size.

2a. In adult outpatients with moderate-severe ulcerative colitis who are naïve to biologic agents, the AGA suggests using infliximab or vedolizumab rather than adalimumab, for induction of remission. *(Conditional recommendation, moderate quality evidence)*

Comment: Patients, particularly those with less severe disease, who place higher value on the convenience of self-administered subcutaneous injection, and a lower value on the relative efficacy of medications, may reasonably chose adalimumab as an alternative.

2b. In adult outpatients with moderate-severe ulcerative colitis who are naïve to biologic agents, the AGA recommends that tofacitinib only be used in the setting of a clinical or registry study. *(No recommendation, knowledge gap)* Comment: Updated FDA recommendations (07/26/2019) on indications for use of tofacitinib in ulcerative colitis recommends its use only after failure of, or intolerance to TNF α antagonists.

In adult outpatients with moderate-severe UC naïve to biologic agents, the guideline panel suggests using infliximab or vedolizumab, rather than adalimumab for induction of remission. Based on updated FDA document on approved indication for tofacitinib use only in patients after failure of, or intolerance to TNF α antagonists, the guideline panel recommends that any use of tofacitinib in biologic-naïve patients with UC be closely monitored in the setting of a clinical or registry study. Currently, both infliximab and vedolizumab are intravenous medications that require infusions, which may be inconvenient to some patients. For these patients, particularly

those with less severe disease, who value the convenience of self-administered injection therapy, adalimumab may be a reasonable alternative option as first-line biologic therapy.

In the head-to-head trials comparing vedolizumab vs. adalimumab in patients with moderate-severe UC (VARSITY), rate of clinical remission was significantly higher in vedolizumab-treated patients vs. adalimumab-treated patients (34.2% vs. 24.3%; RR, 1.41; 95% CI, 1.10-1.81) amongst biologic-naïve patients. For all other comparisons, evidence on comparative efficacy was derived from a network meta-analysis. Network meta-analysis can help assess comparative efficacy of several interventions and synthesize evidence across a network of RCTs, especially if there is weak (or absent) direct evidence.¹⁴ Such indirect comparisons of competing interventions, adjusted by a common control, such as placebo, can partially take account of prognostic characteristics of patients in different trials. The analysis included 15 RCTs with a total of 3747 biologic-naïve patients with moderate-severe UC, treated with infliximab (4 trials, 667 patients), adalimumab (4 trials, 1046 patients), golimumab (2 trials, 586 patients), vedolizumab (3 trials, 630 patients), tofacitinib (2 trials, 520 patients) and ustekinumab (1 trial, 298 patients) were included (see technical review). For this body of evidence for induction therapy, trial design, participant characteristics, interventions, comparators and outcomes for trials of infliximab, adalimumab, golimumab, vedolizumab and ustekinumab were deemed reasonably similar to facilitate indirect comparison. In contrast, trials of tofacitinib were deemed dissimilar since they used a strict rectal bleeding score of zero for outcome assessment (in contrast to other trials which allowed rectal bleeding score of 0 or 1 in outcome assessment). See the technical review and technical review tables 3-5 for full details regarding this analysis. On network meta-analysis, there was moderate confidence in estimates demonstrating the superiority of infliximab over adalimumab (OR, 2.10; 95% CI, 1.16-3.79) (evidence rated down for serious imprecision). It is important to note that in these clinical trials, treatment was not optimized to suggested drug concentrations; it is possible that the efficacy of

infliximab, adalimumab and golimumab may be comparable in patients who achieve adequate drug concentrations given their similar mechanism of action.

2c. In adult outpatients with moderate-severe ulcerative colitis who have previously been exposed to infliximab, particularly those with primary non-response, the AGA suggests using ustekinumab or tofacitinib, rather than vedolizumab or adalimumab for induction of remission. (Conditional recommendation, low quality evidence)

Comment: Patients, particularly those with less severe disease, who place higher value on the potential safety of medications, and a lower value on the relative efficacy of medications, may reasonably chose vedolizumab as an alternative.

In adult outpatients with moderate-severe UC who have previously been exposed to infliximab, especially those with primary non-response, the guideline panel suggests using ustekinumab or tofacitinib rather than vedolizumab or adalimumab for induction of remission.

In the VARSITY trial, ~21% patients had received prior treatment with a TNF α antagonist other than adalimumab. In these patients, there was no significant differences in rates of achieving clinical remission at week 52 (20.3% vs. 16.0%), and the overall body of evidence was deemed to be low quality (rated down for very serious imprecision). A separate network meta-analysis was also performed comparing trials of different agents in patients with moderate-severe UC, who had previously been exposed to TNF α antagonists. The network meta-analysis included 7 RCTs with 1580 patients with prior exposure to TNF α antagonists. Infliximab and golimumab were excluded, since the published studies only included biologic-naïve patients (see technical review Table 5). Of note, all these comparisons were rated down for intransitivity. Prior treatment exposure and response is an important effect modifier. Study level estimates did not report what proportion of patients had exposure to more than one TNF α antagonist and whether patients had exposure to multiple different classes of biologics. There was low confidence in estimates supporting higher efficacy of tofacitinib and ustekinumab over

adalimumab (tofacitinib vs. adalimumab: OR, 11.05; 95% CI, 1.79-68.41; ustekinumab vs. adalimumab: OR, 10.71; 95% CI, 2.01-57.20), and over vedolizumab (tofacitinib vs. vedolizumab: OR, 6.18; 95% CI, 1.00-38.00); ustekinumab vs. vedolizumab: OR, 5.99; 95% CI, 1.13-31.76) for induction of clinical remission in patients with prior exposure to TNF α antagonists.

The guideline panel was explicit in stating these conditional recommendations apply only for patients with prior exposure to infliximab. Given prior evidence supporting superiority of infliximab over adalimumab or golimumab in biologic-naive patients with moderate-severe UC and lack of direct evidence in adalimumab- or golimumab-exposed patients, benefit of switching to vedolizumab, ustekinumab or tofacitinib over infliximab, in patients with prior exposure to adalimumab or golimumab, is uncertain. As noted in AGA's therapeutic drug monitoring guidelines, switching out of class may be reasonable in case of lack of response despite achieving adequate drug concentration.¹⁰

Data for maintenance of remission could not be reliably synthesized using network meta-analysis due to significant differences in clinical trial designs – trials of infliximab and adalimumab were treat-straight through trials, whereas maintenance trials of golimumab, vedolizumab, tofacitinib and ustekinumab re-randomized responders to induction therapy – and lack of stratification of data by prior biologic exposure. However, as with standard clinical practice, once a drug is initiated for induction of remission, if effective, it is typically continued for maintenance of remission as well.

3a. In adult outpatients with active moderate-severe ulcerative colitis, the AGA suggests against using thiopurine monotherapy for INDUCTION of remission. (Conditional recommendation, very low quality of evidence)

3b. In adult outpatients with moderate-severe ulcerative colitis in remission, the AGA suggests using thiopurine monotherapy, rather than no treatment, for MAINTENANCE of remission. (Conditional recommendation low quality of evidence)

The panel suggests against using thiopurine monotherapy for induction of remission in adult outpatients with active moderate-severe UC. However, in patients who have achieved remission (typically induced with corticosteroids), the panel suggests using thiopurine monotherapy rather than no treatment for maintenance of remission.

There were three trials comparing thiopurines vs. placebo and two trials comparing thiopurines vs. 5-aminosalicylates (5-ASA) for inducing corticosteroid free remission.¹⁶⁻²⁰ In four out of five trials, patients were considered corticosteroid-dependent, unable to taper corticosteroids below 10-20mg/day without relapsing. In contrast to modern clinical trials, different disease activity indices were used in these studies, outcome of corticosteroid-free remission was assessed at variable intervals from 4 weeks to 52 weeks, and in patients with active disease started simultaneously on thiopurines and corticosteroids, it was unclear whether remission was induced by corticosteroids or thiopurines. Although thiopurines were associated with a higher rate of corticosteroid-free clinical remission compared to placebo or 5-ASA (RR, 1.25; 95% CI, 1.01-1.56), the overall quality of evidence was deemed very low due to serious risk of bias, imprecision, and indirectness (outcome definition and assessment). Based on the slow onset of action of thiopurines, they are unlikely to be effective as monotherapy for induction of remission in patients with active disease, in the absence of corticosteroids. Hence, based on uncertainty of evidence, the guideline panel opted to suggest against the use of thiopurines for induction of clinical remission in patients with active moderate-severe UC.

For maintenance of remission, four trials comparing thiopurines vs. placebo and three trials comparing thiopurines vs. 5-ASA were included.^{17, 18, 20-24} Maintenance of remission was defined as prevention of relapse following corticosteroid-induced remission (5 trials) or as the

ability to maintain a corticosteroid-free remission in patients on long-standing thiopurine therapy (2 trials), evaluated between 6-18 months. On meta-analysis, thiopurines were more effective than placebo or 5-ASA for prevention of disease relapse (RR, 0.61; 95% CI, 0.49-0.77) among patients with inactive UC (in remission). Quality of evidence was rated as low due to risk of bias and imprecision.

3c. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests against using methotrexate monotherapy, for induction or maintenance of remission.

(Conditional recommendation, low quality evidence)

The guideline panel suggests against the use of methotrexate monotherapy for induction or maintenance of remission for adult outpatients with moderate-severe UC.

Two trials compared methotrexate vs. placebo and one trial compared methotrexate vs. 5-ASA for induction of remission.^{25, 26} In the pivotal METEOR trial, all patients were on 10-40mg/day of corticosteroids with or without active disease.²⁵ The primary outcome was corticosteroid-free remission between weeks 12-30. On meta-analysis, there was no significant difference in rates of inducing remission with methotrexate compared to placebo (RR, 1.31; 95% CI, 0.89-1.94). The quality of evidence was rated as very low due to very serious indirectness (different dosing regimens and modes of administration, variable definition of clinical remission, and inability to truly assess whether remission was induced by corticosteroids or methotrexate), and serious imprecision. For maintenance of remission, two trials compared methotrexate vs. placebo and one compared methotrexate vs. 5-ASA.^{18, 25, 26} Similar to induction, there was no difference between methotrexate and placebo/5-ASA for maintenance of remission (RR, 1.01; 95% CI, 0.79-1.29). The quality of evidence was rated as very low due to serious indirectness, and very serious imprecision.

4a. In adult outpatients with active moderate-severe ulcerative colitis, the AGA suggests using biologic monotherapy (TNF α antagonists, vedolizumab, ustekinumab) rather than thiopurine monotherapy for INDUCTION of remission. (Conditional recommendation, low quality evidence)

4b. In adult outpatients with moderate-severe ulcerative colitis in remission, the AGA makes no recommendation in favor of, or against, using biologic monotherapy (TNF α antagonists, vedolizumab or ustekinumab), rather than thiopurine monotherapy for MAINTENANCE of remission. (No recommendation, knowledge gap)

The panel conditionally suggests use of biologic monotherapy rather than thiopurine for induction of remission. The panel makes no recommendation in favor of or against using biologic monotherapy over thiopurine monotherapy for maintenance of remission.

Evidence was informed by a three-arm clinical trial, UC-SUCCESS comparing infliximab vs. azathioprine vs. combination therapy of infliximab with azathioprine,²⁷ as well as indirect evidence based on trials comparing individual medications with placebo. While UC-SUCCESS was designed as an induction and maintenance trial to evaluate the comparative efficacy of monotherapy vs. combination therapy in biologic-naïve adult outpatients with moderate-severe UC (similar to SONIC for Crohn's disease), the trial was discontinued prematurely by the sponsor before intended enrollment and without completion of the maintenance phase.²⁷ In this trial, there was no difference between infliximab monotherapy and azathioprine monotherapy for achieving corticosteroid-free remission at week 16 (RR, 0.96; 95% CI, 0.53-1.72), albeit based on very low quality evidence. Infliximab monotherapy was superior to thiopurine monotherapy

for achieving endoscopic remission, an important outcome. In addition, based on prior evidence synthesis from PICO 1 and 3 and corresponding recommendations, AGA recommends using biologic agents (TNF α antagonists, vedolizumab or ustekinumab) over placebo (moderate quality evidence), whereas it suggests against using thiopurine monotherapy, for induction of remission (very low quality evidence) in adult outpatients with moderate-severe UC.

Since UC-SUCCESS was terminated prematurely,²⁷ no head-to-head trials informed comparative efficacy of biologic monotherapy vs. thiopurines for maintenance of remission. As in PICO 1 and 3, both strategies have been recommended for maintenance of remission, over no treatment. Recognizing lack of evidence, the panel makes no recommendation in favor of or against biologic monotherapy rather than thiopurine monotherapy, for maintenance of remission. Physician judgment factoring in patients' clinical status, safety profile of different agents and costs and convenience of therapy may be used to inform choice of agents.

5a. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests combining TNF α antagonists, vedolizumab or ustekinumab with thiopurines or methotrexate, rather than biologic monotherapy. (Conditional recommendation, low quality evidence)

Comment: Patients, particularly those with less severe disease, who place higher value on lower risk of adverse events with biologic monotherapy, and lower value on the relative efficacy of combination therapy, may reasonably chose biologic monotherapy.

5b. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests combining TNF α antagonists, vedolizumab or ustekinumab with thiopurines or methotrexate, rather than thiopurine monotherapy. (Conditional recommendation, low quality evidence)

The guideline panel suggests combining TNF α antagonists, vedolizumab or ustekinumab with thiopurines or methotrexate over biologic monotherapy, or thiopurine monotherapy, in adult outpatients with moderate-severe UC. However, in patients particularly those with less severe disease who place higher value on safety and tolerability of pharmacotherapy may reasonably chose biologic monotherapy.

Combination therapy of infliximab with thiopurine compared to infliximab monotherapy was evaluated in UC-SUCCESS, a single three-arm double blind double dummy RCT in biologic-naïve patients with.²⁷ As noted above, this trial was terminated prematurely before planned enrollment and prior to completion of maintenance trial. Patients were randomized to infliximab monotherapy, azathioprine monotherapy, or combined infliximab and azathioprine therapy.²⁷ Combination therapy was more effective for inducing a corticosteroid-free remission at week 16 compared to infliximab monotherapy (RR, 1.78; 95% CI, 1.08-1.94). No trials compared combination therapy of other TNF α antagonists, vedolizumab or ustekinumab with immunomodulators vs. biologic monotherapy. Since adding immunomodulators to biologic agents improves pharmacokinetics of the biologic agent (increasing trough concentration and decreasing immunogenicity), the guideline panel extrapolated this indirect evidence favoring combination therapy of thiopurines and infliximab to other TNF α antagonists, vedolizumab or ustekinumab, particularly in patients with unfavorable pharmacokinetics (more severe disease, higher inflammatory burden, low albumin, higher body mass index), even though the immunogenicity of newer biologic agents may be lower than infliximab.²⁸ Evidence supporting the use of combination therapy vs. immunomodulator monotherapy is also based on UC-SUCCESS.²⁷ In this trial, combination of infliximab and thiopurines was superior to thiopurine monotherapy for achieving corticosteroid-free remission at week 16 (RR, 1.70; 95% CI, 1.04-2.78).

No clinical trial comparing combination therapy with monotherapy for maintenance of remission was identified. Very low quality evidence from a retrospective French study of 82 patients in remission on combination therapy, suggested that continuing combination infliximab and azathioprine was superior to de-escalating to infliximab monotherapy.

Overall, quality of evidence for infliximab-based combination therapy vs. infliximab monotherapy and vs. thiopurine monotherapy for induction of remission was rated as moderate quality (rated down for imprecision). No trials comparing combination therapy of non-infliximab TNF α antagonists (adalimumab, golimumab), vedolizumab or ustekinumab with thiopurines (or methotrexate) vs. biologic monotherapy were identified. Extrapolating from evidence supporting the use of combination of therapy with infliximab, acknowledging lower risk of immunogenicity with newer biologic agents, evidence of other agents was rated as low quality, due to indirectness. Quality of evidence for maintenance of remission was rated as very low due to the observational evidence and serious imprecision.

6. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests early use of biologic agents with or without immunomodulator therapy, rather than gradual step up after failure of 5-aminosalicylates. (Conditional recommendation, very low quality evidence)

Comment: Patients, particularly those with less severe disease, who place higher value on the safety of 5-ASA therapy, and lower value on the efficacy of biologic agents, may reasonably choose gradual step therapy with 5-ASA therapy.

The guideline panel suggests early use of biologics with or without immunomodulator therapy, or tofacitinib rather than gradual step up therapy after failure of 5-aminosalicylates (5-ASA), in patients with moderate-severe disease activity at high risk of colectomy. However, in patients

with less severe disease who place higher value on the safety profile of 5-ASA therapy and lower value on the overall efficacy of biologics may choose to start with 5-ASA therapy.

There were no studies identified that compared a strategy of upfront biologic-based therapy or tofacitinib vs. gradual step-up therapy (introducing biologic-based therapy or tofacitinib only after failure of 5-ASA), or that compared biologic-based therapy vs. 5-ASA-based therapy, in patients with moderate-severe UC at high risk of colectomy. Three studies compared thiopurines vs. 5-ASA, in corticosteroid-exposed patients with UC.^{18, 21, 24} On meta-analysis, thiopurines achieved higher rates of corticosteroid-free remission than 5-ASA. Indirectly extrapolating the data from PICO questions 3 and 4, suggests that biologic-based therapy will likely be more effective than 5-ASA-based therapy. Importantly, 5-ASAs have not been specifically studied in patients with moderate-severe disease activity, and their use is limited to patients with mild-moderate disease activity. Delaying effective treatment to induce remission in patients with moderate-severe UC at high risk of colectomy may be harmful due to ongoing untreated active disease, increasing risk of UC-related complications, hospitalization, colectomy and overall inferior quality of life.

The overall quality of evidence supporting this recommendation was rated as very low due to serious indirectness and imprecision.

7. In adult outpatients with moderate-severe ulcerative colitis who have achieved remission with biologic agents and/or immunomodulators, or tofacitinib, the AGA suggests against continuing 5-aminosalicylates for induction and maintenance of remission. (Conditional recommendation, very low quality evidence)

The guideline panel suggests against continuation of 5-ASA in adult outpatients with moderate-severe UC for induction and maintenance of remission, who have previously failed 5-ASA and have been escalated to a biologic agent and/or immunomodulators or tofacitinib.

In a randomized clinical trial, patients with moderate-severe UC in corticosteroid-free clinical, endoscopic and histologic remission on azathioprine and olsalazine, were randomized to either continuation of azathioprine with olsalazine or de-escalation to azathioprine alone. In two years of follow up, there were no differences in risk of relapse between the two groups (RR, 1.02; 95% CI, 0.77-1.34). There were no studies of systematic withdrawal of 5-ASA in biologic- or tofacitinib-treated patients with moderate-severe UC. Indirect evidence from sub-group analysis of RCTs comparing rates of induction and maintenance of remission in patients who were vs. were not concomitantly on 5-ASA at trial entry was used. On meta-analysis, there was no differences in rates of inducing or maintaining clinical remission in TNF α antagonist- or tofacitinib-treated patients who were vs. were not on concomitant 5-ASA at trial entry (Induction: RR, 0.94; 95% CI, 0.74-1.18| maintenance: RR, 0.92; 95% CI, 0.78-1.09). Of note, these trials did not assess effect of systematic withdrawal of 5-ASAs in biologic- or tofacitinib-treated patients. Overall quality of evidence was rated as low due to imprecision and indirectness of evidence.

This recommendation did not factor in whether continuing 5-ASAs may or may not have a chemoprotective benefit against colorectal cancer in patients with UC. While studies have variably shown an association between 5-ASA use and lower risk of colorectal cancer in patients with UC,^{29, 30} recent data suggests that chronically active disease is a risk factor for colon cancer and that sustained remission is protective regardless of the type of therapy used.³¹

MANAGEMENT OF HOSPITALIZED PATIENTS WITH ACUTE SEVERE ULCERATIVE COLITIS

8. In hospitalized adult patients with acute severe ulcerative colitis, the AGA suggests using intravenous methylprednisolone dose equivalent of 40 to 60mg/d rather than higher dose intravenous corticosteroids. (Conditional recommendation, very low quality evidence)

The guideline panel suggests using intravenous corticosteroids at doses equivalent to 40-60mg/day of methylprednisolone rather than higher doses of intravenous corticosteroids to decrease risk of colectomy in hospitalized adults with ASUC.

Intravenous corticosteroids are the mainstay of management of hospitalized adults with ASUC. There were no head-to-head trials comparing different doses of corticosteroids in hospitalized patients with ASUC. In a systematic review evaluating the risk factors for colectomy in patients with ASUC, Turner and colleagues observed that mean methylprednisolone dose was 68mg/day (range 40-100mg) in hospitalized patients with ASUC; on meta-regression after controlling for baseline disease severity, there was no correlation between corticosteroid dose and risk of colectomy ($R^2 < 0.01$).³² In included trials, different intravenous corticosteroid regimens (different formulations, given once daily vs. multiple times daily vs. as continuous infusion) were used, and none was superior to others. Based on models estimating risk and risk factors for colectomy in patients with ASUC, corticosteroid trials of 3-5 days are suggested; continued use of corticosteroids beyond 7 days has not been shown to be effective in non-responding patients.^{32, 33}

Overall quality of evidence was rated as very low due to observational nature of evidence and indirectness in approach to comparing efficacy.

9. In hospitalized adult patients with acute severe ulcerative colitis without infections, the AGA suggests against adjunctive antibiotics. (conditional recommendation, very low quality of evidence)

The guideline panel suggests against the routine use of adjunctive antibiotics for the treatment of ASUC in patients without gastrointestinal or extra-intestinal infections.

Four RCTs comparing antibiotics vs. no antibiotics/placebo for the treatment of ASUC were identified.³⁴⁻³⁷ Different antibiotics were used, with durations ranging from 5-10 days. On meta-analysis, the addition of antibiotics was not superior to no antibiotics for decreasing short-term risk of colectomy in patients with ASUC (RR, 0.79; 95% CI, 0.46-1.35). After exclusion of one trial of oral vancomycin which was positive but used an insensitive test to exclude concomitant *Clostridium difficile* infection, overall summary estimate for adjunctive antibiotics was close to unity (RR, 0.95; 95% CI, 0.55-1.64).

The quality of evidence was rated very low due to serious risk of bias, serious imprecision and inconsistency given the diverse group of antibiotics used in the studies.

10. In hospitalized adult patients with acute severe ulcerative colitis refractory to intravenous corticosteroids, the AGA suggests using infliximab or cyclosporine. (Conditional recommendation, low quality evidence)

The guideline panel suggests using either infliximab or cyclosporine in hospitalized adult patients with ASUC refractory to a 3-5 day trial of intravenous corticosteroids.

One RCT compared infliximab to placebo in 45 patients with ASUC refractory to intravenous corticosteroids.³⁸ In this trial, infliximab was superior to placebo in decreasing the risk of colectomy within 90 days of hospitalization (RR, 0.44; 95% CI, 0.22-0.87). Patients were given a single dose of 5mg/kg infliximab without further induction or maintenance doses.

Likewise, intravenous cyclosporine (4mg/kg) was compared to placebo in a single small RCT in corticosteroid-refractory patients with ASUC.³⁹ In this trial, there was a trend favoring cyclosporine over placebo (RR, 0.61; 95% CI, 0.18-2.01). In a subsequent RCT, cyclosporine dose 2mg/kg/day was comparable to 4mg/kg/day, with conceivably a superior safety profile with lower dose.⁴⁰ In two head-to-head trials comparing infliximab vs. cyclosporine, there was no significant difference in short-term risk of colectomy between standard dose induction therapy with infliximab vs. cyclosporine in hospitalized patients with corticosteroid-refractory ASUC (RR, 1.00; 95% CI, 0.72-1.40).^{41, 42} Long-term follow-up of these trials also suggest similar findings. Over a median follow-up of 4.5 years of participants in the CYSIF trial, 1- and 5- year colectomy free survival was 70.9% and 61.5% in patients treated with cyclosporine initially and 69.1% and 65.1%, respectively, in patients randomized to infliximab (p=0.97). Importantly, in these trials, randomized treatment was offered for 12-14 weeks, after which treatment decisions were deferred to treating physicians. After the initial randomization period, 1- and 5-year cumulative use of infliximab in cyclosporine-treated patients was 45.7% and 57.1%, respectively; in contrast, only 4 infliximab-treated patients switched to cyclosporine.^{41, 42}

Overall quality of evidence supporting the use of infliximab or cyclosporine over placebo for ASUC refractory to intravenous corticosteroids was rated as moderate quality (imprecision) or low quality (very serious imprecision), respectively. Evidence supporting the comparability of infliximab and cyclosporine decreasing risk of short-term colectomy in hospitalized patients with corticosteroid-refractory ASUC was rated as low quality (open-label trials with high risk of bias, and imprecision).

11. In hospitalized adult patients with acute severe ulcerative colitis, refractory to intravenous corticosteroids, being treated with infliximab, the AGA makes no recommendation on routine use of intensive vs. standard infliximab dosing. (No recommendation, knowledge gap)

The guideline panel makes no recommendation regarding routine use of intensive vs. standard infliximab dosing in adult patients hospitalized with ASUC refractory to intravenous corticosteroids.

There were no RCTs comparing different dosing regimens in patients hospitalized with ASUC. Five observational studies compared outcomes in patients hospitalized with ASUC refractory to intravenous corticosteroids, being treated with different infliximab regimens.⁴³⁻⁴⁶ Intensive dosing regimens included either shortened interval between the infliximab dosing (<2 weeks, dose stacking) and/or induction with higher dose infliximab (10mg/kg) either upfront or at time of dose stacking, without a standard protocol. On meta-analysis of these observational studies, there was no significant difference in the short-term risk of colectomy intensive vs. standard infliximab dosing regimens (RR 1.61; 95% CI, 0.74-3.52). In two studies, upfront induction with higher dose infliximab (10mg/kg) was superior to dose stacking with standard doses (5mg/kg), with lower risk of colectomy (RR, 0.24; 95% CI, 0.08-0.68). Overall, quality of evidence was rated as very low quality due to serious risk of bias, inconsistency and imprecision.

Biologically, patients with corticosteroid-refractory ASUC have a high inflammatory burden that may result in accelerated consumption and fecal wasting of infliximab resulting in low serum concentrations and immunogenicity.⁴⁷ Additionally, given that the drug is albumin bound and many of these patients are malnourished in the setting of ASUC, the systemic drug concentration of biologics may be lower. Hence, it is plausible that intensive regimens may be more effective than standard induction with infliximab. However, this entire body of

observational evidence is confounded by disease severity, wherein patients at intrinsically higher risk of colectomy, or those who have inadequate response to standard induction dose, are treated with more intensive regimens. As a result, given the lack of robust evidence in this setting to guide therapy, this remains a knowledge gap in need of further research to better guide therapy for the ideal induction regimens for infliximab in hospitalized patients with ASUC refractory to intravenous corticosteroids.

SUMMARY

These practice recommendations for the management of moderate-to-severe UC were developed using the GRADE framework and in adherence to the standards established by the Institute of Medicine for the development of trustworthy guidelines^{12, 13}. The goal of this guideline is to promote high quality, high value evidence-based care for patients with moderate-to-severe UC.

Current evidence supports use of infliximab, adalimumab, golimumab, vedolizumab, and tofacitinib for the induction and maintenance of remission in moderate-severe UC. Thiopurine monotherapy should not be used for induction of remission but may be considered for maintenance of remission; in contrast, methotrexate monotherapy, orally or subcutaneously, should not be used for induction or maintenance of remission. Network meta-analysis suggests that infliximab and vedolizumab may be preferred first-line therapy in biologic-naïve patients, rather than standard-dose adalimumab or golimumab, with limited evidence to inform appropriate positioning of tofacitinib. In patients with prior exposure to infliximab, particularly those with primary non-response to induction therapy, vedolizumab or tofacitinib may be preferred over adalimumab or golimumab. Combination therapy of a biologic agent with an immunomodulator is more effective than monotherapy with either agent, though patients, particularly those with less severe disease and those averse to side effects from medications,

may opt for monotherapy. In patients with moderate-severe disease activity, at high risk of colectomy, biologic agents with or without an immunomodulator, or tofacitinib, should be used early rather than gradual step up therapy after failure of 5-aminosalicylates. Patients in remission with biologic agents and/or immunomodulators or tofacitinib after prior failure of 5-ASA, may discontinue 5-aminosalicylates.

Among hospitalized patients with ASUC, after excluding alternative etiologies, intravenous methylprednisolone doses of 40-60mg/day or equivalent are mainstay of therapy. Routine use of adjunctive antibiotics in patients without infections is not recommended. Patients who are refractory to 3-5 days trial of intravenous corticosteroids, who prefer ongoing medical management, may be treated with either infliximab or cyclosporine. In patients being treated with infliximab, no recommendation can be made regarding routine use of intensive vs. standard infliximab dosing.

AREAS FOR FUTURE RESEARCH

The guideline panel identified multiple knowledge gaps and areas for future research in patients with moderate-severe UC. With the increasing number of different drug classes available to treat UC, there is a clear need for identifying biomarkers predictive of response to individual therapies, to facilitate optimal positioning of therapies; in addition, head-to-head trials will directly inform comparative efficacy, and strengthen quality of evidence derived from network meta-analyses. Besides efficacy, safety is an important consideration with different therapies, and different treatment strategies offer distinct risk-benefit profiles. A comprehensive personalization of therapy based on key treatment attributes (efficacy, safety, speed of onset of action, co-interventions, convenience) is warranted to optimally inform shared-decision making. There is limited evidence regarding the utility and duration of combination therapy of biologics and immunomodulators in patients with UC, particularly with newer agents with lower

immunogenicity and with better optimization of biologic agents through therapeutic drug monitoring. Treatment targets with UC are in evolution. It is unclear how well targeting an integrated clinical and biomarker remission compares to endoscopic remission, and whether there may be incremental benefit to treating to a target of histologic remission. Finally, there is considerable paucity of evidence on optimally using existing therapies and novel treatment options for hospitalized patients with ASUC who are refractory to intravenous corticosteroids. The management of these patients, at very high-risk of colectomy in the short-term, also need to be prioritized.

Table 1. Disease severity scoring systems

Truelove and Witts Criteria					
	Mild	Severe	Fulminant		
Stool number/day	<4	>6	>10		
Blood in stool	Intermittent	Frequent	Continuous		
Temperature °C	Normal	>37.5	>37.5		
Pulse beats/minute	Normal	>90	>90		
Hemoglobin	Normal	<75% normal	Transfusion required		
Erythrocyte sedimentation rate mm/hour	≤30	>30	>30		
Colonic features on radiograph	None	Air, edematous wall, thumbprinting	Colonic dilation		
Clinical signs	None	Abdominal tenderness	Abdominal distention and tenderness		
Mayo Score for Ulcerative Colitis					
Variable	Definition	Score	Variable	Definition	Score
Stool pattern	Normal number of daily bowel movements	0 points	Endoscopic findings	Normal/inactive colitis	0 points
	1 to 2 more bowel movements than normal	1 point		Erythema, decreased vascularity	1 point
	3 to 4 more bowel movements than normal	2 points		Friability, marked erythema, erosions	2 points
	5 or more bowel movements than normal	3 points		Ulcerations, severe friability, spontaneous bleeding	3 points
Most severe rectal bleeding of the day	None	0 points	Physician global assessment	Normal	0 points
	Blood streaks seen in the stool < ½ the time	1 point		Mild colitis	1 point
	Blood in most stool	2 points		Moderate colitis	2 points
	Pure blood passed	3 points		Severe colitis	3 points

Table 2. GRADE definitions of quality and certainty of the evidence

Quality Grade	Definition
High	We are very confident that the true effect lies close to the estimate of the effect
Moderate	We are moderately confident in the effect estimate. The true effect is likely to be close to the estimate of effect, but there is a possibility that it is substantially different.
Low	Our confidence in the estimate is limited. The true effect may be substantially different from the estimate of effect.
Very low	We have very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of effect
Evidence gap	Available evidence is insufficient to determine true effect

Table 3. GRADE definitions on strength of recommendation and guide to interpretation

Strength of recommendation	Wording in the guideline	For the patient	For the clinician
Strong	"The AGA recommends..."	Most individuals in this situation would want the recommended course and only a small proportion would not.	Most individuals should receive the recommended course of action. Formal decision aids are not likely to be needed to help individuals make decisions consistent with their values and preferences.
Conditional	"The AGA suggests..."	The majority of individuals in this situation would want the suggested course, but many would not.	Different choices would be appropriate for different patients. Decision aids may be useful in helping individuals in making decisions consistent with their values and preferences. Clinicians should expect to spend more time with patients when working towards a decision.
No recommendation	"The AGA makes no recommendation..."		The confidence in the effect estimate is so low that any effect estimate is speculative at this time.

Table 4. Summary of recommendations of the AGA Clinical Guidelines Committee for the Management of Moderate-to-Severe Ulcerative Colitis

Recommendations	Strength of recommendation	Quality of evidence
<p>1. In adult outpatients with moderate-severe ulcerative colitis, the AGA recommends using infliximab, adalimumab, golimumab, vedolizumab, tofacitinib or ustekinumab over no treatment.</p> <p><i>(Medications are ordered based on year of approval by the US FDA)</i></p>	Strong	Moderate
<p>2a. In adult outpatients with moderate-severe ulcerative colitis who are naïve to biologic agents, the AGA suggests using infliximab or vedolizumab rather than adalimumab, for induction of remission.</p> <p><i>Comment: Patients, particularly those with less severe disease, who place higher value on the convenience of self-administered subcutaneous injection, and a lower value on the relative efficacy of medications, may reasonably chose adalimumab as an alternative.</i></p>	Conditional	Moderate
<p>2b. In adult outpatients with moderate-severe ulcerative colitis who are naïve to biologic agents, the AGA recommends that tofacitinib be only be used in biologic-naïve patients in the setting of a clinical or registry study. (No recommendation, knowledge gap) <i>Comment: Updated FDA recommendations (07/26/2019) on indications for use of tofacitinib in ulcerative</i></p>	No recommendation	Knowledge Gap

colitis recommends its use only after failure of, or intolerance to TNF α antagonists.		
2c. In adult outpatients with moderate-severe ulcerative colitis who have previously been exposed to infliximab, particularly those with primary non-response, the AGA suggests using ustekinumab or tofacitinib, rather than vedolizumab or adalimumab for induction of remission.	Conditional	Low
3a. In adult outpatients with active moderate-severe ulcerative colitis, the AGA suggests against using thiopurine monotherapy for induction of remission.	Conditional	Very low
3b. In adult outpatients with moderate-severe ulcerative colitis in remission, the AGA suggests using thiopurine monotherapy, rather than no treatment, for maintenance of remission.	Conditional	Low
3c. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests against using methotrexate monotherapy, for induction or maintenance of remission.	Conditional	Low
4a. In adult outpatients with active moderate-severe ulcerative colitis, the AGA suggests using biologic monotherapy (TNF-α antagonists, vedolizumab or ustekinumab) or tofacitinib rather than thiopurine monotherapy for induction of remission.	Conditional	Low
4b. In adult outpatients with moderate-severe ulcerative colitis in remission, the AGA makes no recommendation in favor of, or against, using biologic monotherapy or tofacitinib, rather than thiopurine monotherapy for maintenance of remission.	No recommendation	Knowledge gap

<p>5a. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests combining TNFα antagonists, vedolizumab or ustekinumab with thiopurines or methotrexate, rather than biologic monotherapy.</p> <p><i>Comment: Patients, particularly those with less severe disease, who place higher value on the safety of biologic monotherapy, and lower value on the efficacy of combination therapy, may reasonably chose biologic monotherapy.</i></p>	Conditional	Low
<p>5b. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests combining TNF-α antagonists, vedolizumab or ustekinumab with thiopurines or methotrexate rather than thiopurine monotherapy.</p>	Conditional	Low
<p>6. In adult outpatients with moderate-severe ulcerative colitis, the AGA suggests early use of biologic agents with or without immunomodulator therapy, rather than gradual step up after failure of 5-aminosalicylates.</p> <p><i>Comment: Patients, particularly those with less severe disease, who place higher value on the safety of 5-ASA therapy, and lower value on the efficacy of biologic agents or tofacitinib, may reasonably chose gradual step therapy with 5-ASA therapy.</i></p>	Conditional	Very low
<p>7. In adult outpatients with moderate-severe ulcerative colitis who have achieved remission with biologic agents and/or immunomodulators or tofacitinib, the AGA suggests against continuing 5-aminosalicylates for induction and maintenance of</p>	Conditional	Very low

remission.		
8. In hospitalized adult patients with acute severe ulcerative colitis, the AGA suggests using intravenous methylprednisolone dose equivalent of 40 to 60mg/d rather than higher dose intravenous corticosteroids.	Conditional	Very low
9. In hospitalized adult patients with acute severe ulcerative colitis without infections, the AGA suggests against adjunctive antibiotics.	Conditional	Very low
10. In hospitalized adult patients with acute severe ulcerative colitis refractory to intravenous corticosteroids, the AGA suggests using infliximab or cyclosporine.	Conditional	Low
11. In hospitalized adult patients with acute severe ulcerative colitis being treated with infliximab, the AGA makes no recommendation on routine use of intensive vs. standard infliximab dosing.	No recommendation	Knowledge gap

Abbreviations 5-ASA 5-aminosalicylates, AGA American Gastroenterological Association, FDA Food and Drug Administration, TNF tumor necrosis factor, UC ulcerative colitis, US United States

REFERENCES

1. Singh S, Feuerstein JD, Binion DG, et al. American Gastroenterological Association technical review on the management of mild-moderate ulcerative colitis. *Gastroenterology* 2018.
2. Fumery M, Singh S, Dulai PS, et al. Natural History of Adult Ulcerative Colitis in Population-based Cohorts: A Systematic Review. *Clin Gastroenterol Hepatol* 2017;16:343-356 e3.
3. Fumery M, Singh S, Dulai PS, et al. Natural History of Adult Ulcerative Colitis in Population-based Cohorts: A Systematic Review. *Clin Gastroenterol Hepatol* 2018;16:343-356 e3.
4. Narula N, Marshall JK, Colombel JF, et al. Systematic Review and Meta-Analysis: Infliximab or Cyclosporine as Rescue Therapy in Patients With Severe Ulcerative Colitis Refractory to Steroids. *American Journal of Gastroenterology* 2016;111:477-91.
5. Dassopoulos T, Cohen RD, Scherl EJ, et al. Ulcerative Colitis Care Pathway. *Gastroenterology* 2015;149:238-45.
6. Schroeder KW, Tremaine WJ, Ilstrup DM. Coated oral 5-aminosalicylic acid therapy for mildly to moderately active ulcerative colitis. A randomized study. *N Engl J Med* 1987;317:1625-9.
7. Truelove SC, Witts LJ. Cortisone in ulcerative colitis; final report on a therapeutic trial. *Br Med J* 1955;2:1041-8.
8. Ungaro R, Mehandru S, Allen PB, et al. Ulcerative colitis. *Lancet* 2017;389:1756-1770.
9. Ko CW, Singh S, Feuerstein JD, et al. American Gastroenterological Association Institute Guideline on the Management of Mild-to-Moderate Ulcerative Colitis. *Gastroenterology* 2018.
10. Feuerstein JD, Nguyen GC, Kupfer SS, et al. American Gastroenterological Association Institute Guideline on Therapeutic Drug Monitoring in Inflammatory Bowel Disease. *Gastroenterology* 2017;153:827-834.
11. American Gastroenterological Association. AGA Institute clinical practice guideline development process. Volume 2018.
12. Sultan S, Falck-Ytter Y, Inadomi JM. The AGA institute process for developing clinical practice guidelines part one: grading the evidence. *Clin Gastroenterol Hepatol* 2013;11:329-32.
13. Institute of Medicine. *Clinical Practice Guidelines We Can Trust*. Washington, DC.: National Academies Press, 2011.
14. Cipriani A, Higgins JP, Geddes JR, et al. Conceptual and technical challenges in network meta-analysis. *Annals of internal medicine* 2013;159:130-137.
15. Singh S, George J, Boland BS, et al. Primary non-response to tumor necrosis factor antagonists is associated with inferior response to second-line biologics in patients with inflammatory bowel diseases: a systematic review and meta-analysis. *Journal of Crohn's and Colitis* 2018;12:635-643.
16. Caprilli R, Carratu R, Babbini M. Double-blind comparison of the effectiveness of azathioprine and sulfasalazine in idiopathic proctocolitis. Preliminary report. *Am J Dig Dis* 1975;20:115-20.
17. Jewell DP, Truelove SC. Azathioprine in ulcerative colitis: final report on controlled therapeutic trial. *British Medical Journal* 1974;4:627-30.
18. Mate-Jimenez J, Hermida C, Cantero-Perona J, et al. 6-Mercaptopurine or methotrexate added to prednisone induces and maintains remission in steroid-dependent inflammatory bowel disease. *European Journal of Gastroenterology and Hepatology* 2000;12:1227-1233.

19. Rosenberg JL, Wall AJ, Levin B, et al. A controlled trial of azathioprine in the management of chronic ulcerative colitis. *Gastroenterology* 1975;69:96-9.
20. Sood A, Midha V, Sood N, et al. Role of azathioprine in severe ulcerative colitis: one-year, placebo-controlled, randomized trial. *Indian Journal of Gastroenterology* 2000;19:14-6.
21. Ardizzone S, Maconi G, Russo A, et al. Randomised controlled trial of azathioprine and 5-aminosalicylic acid for treatment of steroid dependent ulcerative colitis. *Gut* 2006;55:47-53.
22. Hawthorne AB, Logan RF, Hawkey CJ, et al. Randomised controlled trial of azathioprine withdrawal in ulcerative colitis. *BMJ* 1992;305:20-2.
23. Sood A, Kaushal V, Midha V, et al. The beneficial effect of azathioprine on maintenance of remission in severe ulcerative colitis. *Journal of gastroenterology*. Volume 37, 2002:270-274.
24. Sood A, Midha V, Sood N, et al. Azathioprine versus sulfasalazine in maintenance of remission in severe ulcerative colitis. *Indian journal of gastroenterology*. Volume 22, 2003:79-81.
25. Herfarth H, Barnes EL, Valentine JF, et al. Methotrexate is not superior to placebo in maintaining steroid-free response or remission in ulcerative colitis. *Gastroenterology* 2018.
26. Oren R, Arber N, Odes S, et al. Methotrexate in chronic active ulcerative colitis: a double-blind, randomized, Israeli multicenter trial. *Gastroenterology* 1996;110:1416-21.
27. Panaccione R, Ghosh S, Middleton S, et al. Combination therapy with infliximab and azathioprine is superior to monotherapy with either agent in ulcerative colitis. *Gastroenterology*. Volume 146, 2014:392-400.e3.
28. Vermeire S, Gils A, Accossato P, et al. Immunogenicity of biologics in inflammatory bowel disease. *Therapeutic advances in gastroenterology* 2018;11:1756283X17750355.
29. Nguyen GC, Gulamhusein A, Bernstein CN. 5-aminosalicylic acid is not protective against colorectal cancer in inflammatory bowel disease: a meta-analysis of non-referral populations. *The American journal of gastroenterology* 2012;107:1298.
30. O'connor A, Packey CD, Akbari M, et al. Mesalamine, but not sulfasalazine, reduces the risk of colorectal neoplasia in patients with inflammatory bowel disease: an agent-specific systematic review and meta-analysis. *Inflammatory bowel diseases* 2015;21:2562-2569.
31. Rubin DT, Huo D, Kinnucan JA, et al. Inflammation is an independent risk factor for colonic neoplasia in patients with ulcerative colitis: a case-control study. *Clinical gastroenterology and hepatology* 2013;11:1601-1608. e4.
32. Turner D, Walsh CM, Steinhart AH, et al. Response to Corticosteroids in Severe Ulcerative Colitis: A Systematic Review of the Literature and a Meta-Regression. *Clinical Gastroenterology and Hepatology* 2007;5:103-110.
33. Seo M, Okada M, Yao T, et al. Evaluation of the clinical course of acute attacks in patients with ulcerative colitis through the use of an activity index. *J Gastroenterol* 2002;37:29-34.
34. Chapman RW, Selby WS, Jewell DP. Controlled trial of intravenous metronidazole as an adjunct to corticosteroids in severe ulcerative colitis. *Gut* 1986;27:1210-2.
35. Dickinson RJ, O'Connor HJ, Pinder I. Double blind controlled trial of oral vancomycin as adjunctive treatment in acute exacerbations of idiopathic colitis. *Gut* 1985;26:1380-1384.
36. Mantzaris GJ, Hatzis A, Kontogiannis P, et al. Intravenous tobramycin and metronidazole as an adjunct to corticosteroids in acute, severe ulcerative colitis. *American Journal of Gastroenterology* 1994;89:43-6.

37. Mantzaris GJ, Petraki K, Archavlis E, et al. A prospective randomized controlled trial of intravenous ciprofloxacin as an adjunct to corticosteroids in acute, severe ulcerative colitis. *Scandinavian Journal of Gastroenterology* 2001;36:971-974.
38. Jarnerot G, Hertvig E, Friis-Liby I, et al. Infliximab as rescue therapy in severe to moderately severe ulcerative colitis: a randomized, placebo-controlled study. *Gastroenterology* 2005;128:1805-11.
39. Lichtiger S, Present DH, Kornbluth A, et al. Cyclosporine in severe ulcerative colitis refractory to steroid therapy. *New England Journal of Medicine* 1994;330:1841-5.
40. Van Assche G, D'Haens G, Noman M, et al. Randomized, double-blind comparison of 4 mg/kg versus 2 mg/kg intravenous cyclosporine in severe ulcerative colitis. *Gastroenterology* 2003;125:1025-1031.
41. Laharie D, Bourreille A, Branche J, et al. Ciclosporin versus infliximab in patients with severe ulcerative colitis refractory to intravenous steroids: a parallel, open-label randomised controlled trial. *Lancet* 2012;380:1909-15.
42. Williams JG, Alam MF, Alrubaiy L, et al. Infliximab versus ciclosporin for steroid-resistant acute severe ulcerative colitis (CONSTRUCT): a mixed methods, open-label, pragmatic randomised trial. *The Lancet. Gastroenterology & Hepatology* 2016;1:15-24.
43. Choy MC, Seah D, Gorelik A, et al. Predicting response after infliximab salvage in acute severe ulcerative colitis. *J Gastroenterol Hepatol* 2018;33:1347-1352.
44. Gibson DJ, Heetun ZS, Redmond CE, et al. An accelerated infliximab induction regimen reduces the need for early colectomy in patients with acute severe ulcerative colitis. *Clin Gastroenterol Hepatol* 2015;13:330-335 e1.
45. Govani SM WA, Stidham RW, Higgins PDR, Hardiman K. Accelerated Dosing of Infliximab Prevents Colectomy Within 90 Days in Only Half of Patients With Severe Ulcerative Colitis. *Gastroenterology* 2016;150:S106.
46. Shah SC, Naymagon S, Cohen BL, et al. There is Significant Practice Pattern Variability in the Management of the Hospitalized Ulcerative Colitis Patient at a Tertiary Care and IBD Referral Center. *J Clin Gastroenterol* 2018;52:333-338.
47. Brandse JF, van den Brink GR, Wildenberg ME, et al. Loss of Infliximab Into Feces Is Associated With Lack of Response to Therapy in Patients With Severe Ulcerative Colitis. *Gastroenterology* 2015;149:350-5 e2.